


RLUH Community AKI Guideline



****Discuss with renal AKI team / on call renal Registrar if required and appropriate**

RLUH Community AKI Guideline

AKI Definition: New abnormal renal function; serum creatinine rise ≥ 26 $\mu\text{mol/L}$ or 1.5 times from baseline or oliguria (<0.5 mL/kg/hour for > 6 consecutive hours)

**Table 1 : Definition/Staging system for acute kidney injury		
Stage	Serum creatinine (SCr) criteria	Urine output criteria
1	increase ≥ 26 $\mu\text{mol/L}$ within 48hrs or increase ≥ 1.5 to 1.9 X reference SCr	<0.5 mL/kg/hr for > 6 consecutive hrs
2	increase ≥ 2 to 2.9 X reference SCr ¹	<0.5 mL/kg/ hr for > 12 hrs
3	increase ≥ 3 X reference SCr or increase 354 $\mu\text{mol/L}$ or commenced on renal replacement therapy (RRT) irrespective of stage	<0.3 mL/kg/ hr for > 24 hrs or anuria for 12 hrs

KDIGO staging system for acute kidney injury; ¹reference SCr = baseline SCr

²AKI risk Factors:

Hypotension	Known CKD
Hypovolaemia	Diabetes
Sepsis	Heart Failure
Oliguria	Liver disease
Urological obstruction	Age ≥ 65

Cognitive impairment that may limit access to fluids
 Drugs with nephrotoxic potential (NSAID/ ACEI/ARB/Diuretics/Opiates/ Iodinated contrast agents)

³ Hospital Referral for admission:

Suspected vasculitis/ acute nephritis	Oliguric AKI, not improving
Sepsis with hypotension and AKI	AKI with Hyperkalaemia (Serum K > 6 mmol/L)
AKI requiring Intravenous hydration	
Rapidly progressive AKI (progressive higher stages of AKI in 24-48 hours)	

⁴**Renal screen:** ANCA, Anti-GBM Ab, Serum Immunoglobulin & paraprotein, urine Bence Jones Protein, Urine albumin creatinine ratio

*****CONTACT INFORMATION*****		
Main Kidney Unit	Kidney Referral Contacts	On Call Contacts
Royal Liverpool Uni. Hospital	Fax: 0151 706 5439	AKI specialist nurse bleep 5150 / 4620 Renal SpR at RLUH on bleep through RLUH switch board (0151 706 2000): 24 hours

¹MEWS: Modified Early Warning Score (systolic blood pressure, heart rate, respiratory rate, body temperature, level of consciousness) ³SBP- systolic blood press; S.Cr= serum creatinine

*Oliguria = urine output less than 0.5 ml/kg/hour; CKD (adults with eGFR less than 60 ml/min/1.73 m²)

'NICE' Clinical Guideline 50: <http://guidance.nice.org.uk/CG50/QuickRefGuide/pdf/English>

NICE Acute kidney injury (CG169) : <http://guidance.nice.org.uk/cg169>

©This AKI guideline may not be copied or reproduced in any form without the prior written permission of Renal Unit, Royal Liverpool University Hospital NHS trust.